



MEETING ABSTRACT

Open Access

High-pass-filter cut-offs optimization of the filter-based fatigue index during dynamic contractions

Jungyoon Kim, Sunwoo Park, Youngho Kim*

From 4th Congress of the International Foot and Ankle Biomechanics (i-FAB) Community
Busan, Korea. 8-11 April 2014

Background

Traditional muscle fatigue indices (F_{mean} and F_{med}) have a relatively low sensitivity under dynamic exercise conditions. Various muscle fatigue indices have been attempted to overcome this problem [1,2]. However, their methods

required large amounts of computation and had limitations in time-frequency resolution [3]. Kim et al. introduced the use of a filter-based fatigue index (FI_{hlopt}), which the ratio of high-frequency to low-frequency components of EMG power [3]. In this study, we optimized the cut-offs of the




Figure 1 Linear representations of peak power output versus various EMG indices for all subjects. Peak power output versus FI_{hlopt} of (a) RF muscle; (b) VM muscle; (c) VL muscle; (d) BB muscle

* Correspondence: younghokim@yonsei.ac.kr
Department of Biomedical Engineering and Institute of Medical Engineering,
Yonsei University, Wonju, Gangwon, 220-710, Korea

high-pass-filter (HPF) to maximize the correlation coefficient between the peak power and the FI_{hlrOPT} in different muscles, and then to determine the frequency bandwidth of our fatigue index.

Methods

Forty-one healthy males were recruited for this study. Twenty-seven subjects performed knee extension/flexion exercises and fourteen subjects performed elbow flexion/extension exercises on an isokinetic dynamometer (Biodex System 3, Biodex Medical Systems, NY, USA). 10 repetition maximum was used for fatigue exercises. The experimental protocol was 5 sets of 10 knee extensions and elbow flexion with 2 minutes of rest between sets. EMG signals were obtained from rectus femoris (RF), vastus medialis (VM), vastus lateralis (VL) and biceps brachii (BB) muscles using the Noraxon EMG System (MyoSystem 1200, Noraxon Inc., AZ, USA). EMG signals, as well as biomechanical signals (angle, angular velocity and torque) were simultaneously recorded with a sampling rate of 1kHz. FI_{hlrOPT} was calculated and cut-offs of HPF were optimized to maximize the correlation between the peak power and FI_{hlrOPT} .

Results

Optimized cut-offs of RF, VM, VL and BB were similar (353.3 ± 49.5 Hz, 343.9 ± 34.2 Hz, 353.7 ± 36.1 Hz and 362.3 ± 28.2 Hz). RF, VM, VL and BB muscles showed good correlation with joint power (correlation coefficient was 0.81 ± 0.08 , 0.56 ± 0.23 , 0.52 ± 0.24 and 0.72 ± 0.08).

Discussion

Similar to previous study, cut-offs of HPF of muscles were about 350Hz (RF: 360Hz in previous study). Similar cut-offs of HPF in different muscles showed the possibility of general cut-off for muscle fatigue estimation. Mills showed that the compound muscle action potential spectrum did not change during fatigue above 200Hz [4]. Thus, the high frequency power decreased during fatigue because of the reduced motor unit activation. The low frequency power increased during fatigue because of the elevated motor unit action potential areas. FI_{hlrOPT} could reflect the decrease in the peak power during fatigue by these reasons.

Acknowledgement

This research project was also supported by the Sports Promotion Fund of Seoul Olympic Sports Promotion Foundation from Ministry of Culture, Sports and Tourism (s07201212022012).

Published: 8 April 2014

References

1. Karlsson S, Yu J, Akay M: Time-frequency analysis of myoelectric signals during dynamic contractions. *IEEE Trans. Biomed. Eng* 2000, **47**:229-238.
2. Dimitrov GV, Arabadzhiev TI, Mileva KN, Bowtell JL, Crichton N, Dimitrova NA: Muscle fatigue during dynamic contractions assessed by new spectral indices. *Med. Sci. Sports Exerc* 2006, **38**:1971-1979.

3. Kim J, Park S, Ahn S, Kim Y: A Novel Approach of Defining Fatigue Indices with sEMG Power during Isotonic Contractions. *J. Precis. Eng. Manuf* 2012, **13**:977-983.
4. Mills KR: Power spectral analysis of electromyogram and compounded muscle action potential during muscle fatigue and recovery. *J. Physiol* 1982, **326**:401-409.

doi:10.1186/1757-1146-7-S1-A129

Cite this article as: Kim et al.: High-pass-filter cut-offs optimization of the filter-based fatigue index during dynamic contractions. *Journal of Foot and Ankle Research* 2014 **7**(Suppl 1):A129.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

